

Nitric oxide and hydrogen sulfide share regulatory functions in higher plant events

FRANCISCO J. CORPAS^{*}; SALVADOR GONZÁLEZ-GORDO; MARTA RODRÍGUEZ-RUIZ; MARÍA A. MUÑOZ-VARGAS; JOSÉ M. PALMA

Group of Antioxidants, Free Radicals and Nitric Oxide in Biotechnology, Food and Agriculture, Department of Biochemistry, Cell and Molecular Biology of Plants, Estación Experimental del Zaidín (Spanish National Research Council, CSIC), Granada, 18008, Spain

Key words: Ascorbate peroxidase, Hydrogen sulfide, Nitric oxide, Persulfidation, S-nitrosation, Signaling

Abstract: Nitric oxide (NO) and hydrogen sulfide (H₂S) are two molecules that share signaling properties in plant and animal cells. NO and H₂S originate two families of derived molecules designated reactive nitrogen and sulfur species (RNS and RSS, respectively). These molecules are responsible for certain protein regulatory processes through posttranslational modifications (PTMs), being the most remarkable S-nitrosation and persulfidation, which affect the thiol group of cysteine residues. NO and H₂S can also exert regulatory functions due to their interaction through the iron present in proteins that contain heme groups or iron-sulfur clusters, as reported mainly in animal cells. However, the available information in plant cells is still very limited thus far. In higher plants, NO and H₂S are involved in a myriad of physiological events from seed germination to fruit ripening, but also the mechanism of response to biotic and abiotic stress conditions. This viewpoint manuscript highlights the functional regulatory parallelism of these two molecules which also interact with the metabolism of reactive oxygen species (ROS) in plant cells.

Brief Historical Perspective

Nitric oxide (NO) and hydrogen sulfide (H₂S) are two gaseous molecules that were initially considered dangerous because they were associated with some detrimental effects on animal and plant cells. However, this perspective underwent a drastic change of mind when it was found that these molecules were endogenously generated in animal cells (Kolluru *et al.*, 2013). There was a gap of about 10 years between the initial research works that described the signaling functions of either NO or H₂S in living organisms. Accordingly, key research on NO, published in 1987 (Palmer *et al.*, 1987), and on H₂S in 1996 (Abe and Kimura, 1996) in animal systems provided the first pieces of evidence showing that these molecules exerted diverse signaling roles in the cardiovascular and nervous systems, respectively. Years later, plant biologists also found that these molecules were also endogenously generated in plant cells where they are involved in almost all of the stages of plant development including seed germination, root development, plant growth, stomata movement, senescence,

flowering and fruit ripening (Leshem *et al.*, 1998; Lamattina *et al.*, 2003; Simontacchi *et al.*, 2004; Corpas *et al.*, 2004; Corpas *et al.*, 2006; Corpas *et al.*, 2008; Zhou *et al.*, 2018; Chen *et al.*, 2019; González-Gordo *et al.*, 2019; Mukherjee and Corpas, 2020; Zuccarelli *et al.*, 2021). And both compounds were also linked to the mechanisms of response against adverse environmental conditions triggered by either abiotic or biotic agents (Corpas, 2019; Kharbech *et al.*, 2020; Iqbal *et al.*, 2021). Fig. 1 illustrates the key functions in which NO and H₂S have been shown to participate in higher plants.

Chemistry and Biochemistry of NO and H₂S

Although NO and H₂S are very simple molecules, their (bio) chemistry is more complex than it could be thought (Stamler *et al.*, 1992; McCleverty, 2004; Hughes, 2008; Kabil and Banerjee, 2010; Filipovic *et al.*, 2018; González-Gordo *et al.*, 2020). NO is a colourless gas that belongs to the free radical-type molecules because it has an unpaired electron in the π orbital of the nitrogen atom, what is usually indicated with a dot in the chemical formula (*NO). Some of the NO and H₂S physical and chemical properties are: (i) Solubility of NO is 1.9 mM in aqueous solutions at 1 atm pressure, whereas the solubility of H₂S is 100 mM at the same pressure; (ii) Their *in vivo* lifetime is relatively short,

^{*}Address correspondence to: Francisco J. Corpas, javier.corpas@eez.csic.es
Received: 29 April 2021; Accepted: 01 June 2021






FIGURE 1. Main processes where both nitric oxide (NO) and hydrogen sulfide (H_2S) are involved in higher plants.

less than 10 seconds for NO and between second to few minutes for H_2S . Therefore, they can trigger regulatory functions in cellular loci far from their production sites; (iii) NO and H_2S are lipophilic molecules and they can diffuse across cell membranes; and, (iv) Both molecules can react with thiol groups from peptides and proteins affecting the function of the target molecules.

Likewise, NO and H_2S can also interact with iron containing-proteins, where the metal is present as either heme group or as part of the iron-sulfur cluster. Thus, there are multiple examples in higher plants where either NO and H_2S , or both, can modulate, through their interaction with the cysteine thiol groups, the functions of proteins such as cytochrome *c* oxidase, catalase, Fe-superoxide dismutase, ascorbate peroxidase, ferredoxin(Fd)-NADP reductase, glutaredoxin, Fd-dependent glutamine:2-oxyoglutarate aminotransferase (Fd-GOGAT) or phytooglobins, (Ramirez *et al.*, 2011; Aroca *et al.*, 2017; Bahmani *et al.*, 2019; Palma *et al.*, 2020; Niu *et al.*, 2019; Corpas *et al.*, 2021). These proteins are involved in essential plant processes including photosynthesis, respiration, antioxidant system, nitrogen and sulfur assimilation, which remarks the physiological relevance of these signaling molecules. However, in higher plants, the available information about the direct interaction of NO and H_2S with the iron side of protein is still scarce. In addition to reactions that can originate the respective families of molecules derived from nitric oxide and hydrogen sulfide (RNS and RSS, respectively), NO and H_2S can interfere with the biosynthesis of each other and also produce novel species through their chemical interaction, thereby expanding the network of interactions that can affect to macromolecules (Kolluru *et al.*, 2013; Scuffi *et al.*, 2014; Kolluru *et al.*, 2015; Hancock and Whiteman, 2016; Iqbal *et al.*, 2021).

S-nitrosation and Persulfidation: Two Protein PTMs that Exert Redox Control of Thiol Groups

The amino acid cysteine (Cys) can play relevant roles in proteins such as a structural function through disulfide bonds, but it could also have implications on redox reactions by means of its thiol group ($-SH$). Thus, depending on the conditions surrounding the thiol group in the protein, Cys can be found in its anionic form, designated as thiolate (RS^-), which is a stronger nucleophilic

agent than its protonated form (Netto *et al.*, 2007). Fig. 2A depicts the different oxidation states of sulfur which range from thiol (-2) to sulfonic acid ($+4$). Among these states, NO or H_2S can interact with the thiolate form through either S-nitrosation or persulfidation (Fig. 2B), also known previously as protein S-nitrosylation and S-sulfhydration, respectively (Aroca *et al.*, 2018; Wolhuter *et al.*, 2018; Corpas *et al.*, 2019, Corpas *et al.*, 2021). For that reason, Cys is considered as a redox switch in the protein metabolism because it is the main target of these two PTMs, and this could affect significantly the biological activity of the corresponding protein, either positively or negatively.

Ascorbate Peroxidase (APX) in Plant Cells: A Case Study of NO and H_2S Target

In-plant systems, the number of identified proteins that undergo PTMs mediated by either NO or H_2S has progressively increased thanks to the efforts of many researchers focused on this biochemical area (Lindermayr *et al.*, 2005; Tanou *et al.*, 2009; Fares *et al.*, 2011; Begara-Morales *et al.*, 2013; Kato *et al.*, 2013; Chen *et al.*, 2014; Aroca *et al.*, 2015; Liu *et al.*, 2019). Moreover, the analyses of these modified proteins have revealed that many of them can be the simultaneous target of both PTMs and, by *in vitro* assays, it has been also proven the relevance of these two regulatory molecules to modulate the biological activity of the affected proteins (Muñoz-Vargas *et al.*, 2018, 2020; Palma *et al.*, 2020; Corpas *et al.*, 2021). Among the different plant proteomic studies focused on the identification of the potential targets of PTMs mediated by either NO or H_2S , it has been found that ascorbate peroxidase (APX) is one of those shared targets.




FIGURE 2. (A) Oxidation states of sulfur (S) in proteins from thiol (-2) to sulfonic acid ($+4$) forms. Under cellular oxidant conditions, the oxidation from sulfenic acid becomes irreversible. The numbers in parenthesis represent the different oxidation states of S in the protein. (B) Protein thiol modifications mediated by NO (S-nitrosation) and H_2S (persulfidation).

APX is a key antioxidant enzyme that is part of the ascorbate-glutathione cycle, which is an essential system to modulate the mechanism of response against (a)biotic stress environmental conditions (Shigeoka *et al.*, 2002; Asada, 2006; Maruta and Ishikawa, 2018). APX is a hemoprotein that controls the cellular level of hydrogen peroxide (H₂O₂) according to the following reaction:



This enzyme system is composed of different isozymes located in almost all subcellular compartments including cytosol, chloroplasts, mitochondria and peroxisomes (Asada, 1992; Yamaguchi *et al.*, 1995; Bunkelmann and Trelease, 1996; Jiménez *et al.*, 1998; Yoshimura *et al.*, 1999; Maruta *et al.*, 2016; Chin *et al.*, 2019). This molecular and location diversity suggests the great relevance of APX in cell signaling under physiological and stressful conditions, which is consolidated by its regulation through both S-nitrosation and persulfidation, as indicated above. Furthermore, it was identified by mass spectrometric analyses that the Cys32 is the residue that underwent S-nitrosation and persulfidation (Begara-Morales *et al.*, 2014; Yang *et al.*, 2015; Aroca *et al.*, 2015) and, in both cases, the APX activity was positively regulated. This mimicking biochemical regulation provides a clear connection between NO and H₂S with the metabolism of reactive oxygen species (ROS) (Rodríguez-Ruiz *et al.*, 2017), thus indicating the biochemical link among all these families of molecules.

Conclusions and Future Perspectives

The previous perception of NO and H₂S as harmful molecules to plant cells has drastically changed and, nowadays, they are key signal molecules that regulate a myriad of biochemical and physiological processes. These two gases have also families of derived molecules designated as reactive nitrogen and sulfur species (RNS and RSS, respectively). They have a wide range of biochemical implications, being S-nitrosation and persulfidation two representative examples of their cellular relevance which compete molecularly to modulate protein functions through the reaction with the thiol group of cysteines. Additionally, NO and H₂S have been started to be considered as molecules with biotechnological properties since when they are applied exogenously they can exert beneficial effects on crops (Corpas *et al.*, 2019; Corpas and Palma, 2020; Corpas *et al.*, 2020). Therefore, it could be concluded that NO and H₂S behave as competing but mimicking molecules that they can reinforce each other in their signaling properties.

Acknowledgement: SG-G and MAM-V acknowledge respective “Formación de Personal Investigador” contracts from the Ministry of Science and Innovation, Spain.

Authors’ Contribution: FJC conceived the work and wrote the first draft. SGG, MRR, MAMV and JMP critically reviewed the draft. All authors contributed to drafting the work, revised the final manuscript, and approved submission.

Funding Statement: FJC and JMP research is supported by a European Regional Development Fund cofinanced grant from the Spanish Ministry of Science and Innovation

(PID2019-103924GB-I00), the Plan Andaluz de Investigación, Desarrollo e Innovación (PAIDI 2020) (P18-FR-1359) and Junta de Andalucía (Group BIO192), Spain.

Conflicts of Interest: The authors declare that they have no conflicts of interest to report regarding the present study.

References

- Abe K, Kimura H (1996). The possible role of hydrogen sulfide as an endogenous neuromodulator. *Journal of Neuroscience* **16**: 1066–1071.
- Aroca Á, Serna A, Gotor C, Romero LC (2015). S-sulfhydration: A cysteine posttranslational modification in plant systems. *Plant Physiology* **168**: 334–342.
- Aroca A, Benito JM, Gotor C, Romero LC (2017). Persulfidation proteome reveals the regulation of protein function by hydrogen sulfide in diverse biological processes in Arabidopsis. *Journal of Experimental Botany* **68**: 4915–4927.
- Aroca A, Gotor C, Romero LC (2018). Hydrogen sulfide signaling in plants: Emerging roles of protein persulfidation. *Frontiers in Plant Science* **9**: 1369.
- Asada K (1992). Ascorbate peroxidase: A hydrogen peroxide scavenging enzyme in plants. *Physiologia Plantarum* **85**: 235–241.
- Asada K (2006). Production and scavenging of reactive oxygen species in chloroplasts and their functions. *Plant Physiology* **141**: 391–396.
- Bahmani R, Kim D, Na J, Hwang S (2019). Expression of the tobacco non-symbiotic class 1 hemoglobin gene Hb1 reduces cadmium levels by modulating Cd transporter expression through decreasing nitric oxide and ROS level in Arabidopsis. *Frontiers in Plant Science* **10**: 201.
- Begara-Morales JC, López-Jaramillo FJ, Sánchez-Calvo B, Carreras A, Ortega-Muñoz M, Santoyo-González F, Corpas FJ, Barroso JB (2013). Vinyl sulfone silica: Application of an open preactivated support to the study of transnitrosylation of plant proteins by S-nitrosoglutathione. *BMC Plant Biology* **13**: 61.
- Begara-Morales JC, Sánchez-Calvo B, Chaki M, Valderrama R, Mata-Pérez C, López-Jaramillo J, Padilla MN, Carreras A, Corpas FJ, Barroso JB (2014). Dual regulation of cytosolic ascorbate peroxidase (APX) by tyrosine nitration and S-nitrosylation. *Journal of Experimental Botany* **65**: 527–538.
- Bunkelmann JR, Trelease RN (1996). Ascorbate peroxidase. A prominent membrane protein in oilseed glyoxysomes. *Plant Physiology* **110**: 589–598.
- Chen J, Liu TW, Hu WJ, Simon M, Wang WH, Chen J, Liu X, Zheng HL (2014). Comparative proteomic analysis of differentially expressed proteins induced by hydrogen sulfide in *Spinacia oleracea* leaves. *PLoS One* **9**: e105400.
- Chen Z, Huang Y, Yang W, Chang G, Li P, Wei J, Yuan X, Huang J, Hu X (2019). The hydrogen sulfide signal enhances seed germination tolerance to high temperatures by retaining nuclear COP1 for HY5 degradation. *Plant Science* **285**: 34–43.
- Chin DC, Senthil Kumar R, Suen CS, Chien CY, Hwang MJ, Hsu CH, Xuhan X, Lai ZX, Yeh KW (2019). Plant cytosolic ascorbate peroxidase with dual catalytic activity modulates abiotic stress tolerances. *iScience* **16**: 31–49.
- Corpas FJ (2019). Hydrogen sulfide: A new warrior against abiotic stress. *Trends in Plant Science* **24**: 983–988.
- Corpas FJ, Barroso JB, Carreras A, Quirós M, León AM, Romero-Puertas MC, Esteban FJ, Valderrama R, Palma JM, Sandalio

- LM, Gómez M, del Río LA (2004). Cellular and subcellular localization of endogenous nitric oxide in young and senescent pea plants. *Plant Physiology* **136**: 2722–2733.
- Corpas FJ, Barroso JB, Carreras A, Valderrama R, Palma JM, León AM, Sandalio LM, del Río LA (2006). Constitutive arginine-dependent nitric oxide synthase activity in different organs of pea seedlings during plant development. *Planta* **224**: 246–254.
- Corpas FJ, Chaki M, Fernández-Ocaña A, Valderrama R, Palma JM, Carreras A, Begara-Morales JC, Airaki M, del Río LA, Barroso JB (2008). Metabolism of reactive nitrogen species in pea plants under abiotic stress conditions. *Plant Cell and Physiology* **49**: 1711–1722.
- Corpas FJ, González-Gordo S, Palma JM (2020). Nitric oxide: A radical molecule with potential biotechnological applications in fruit ripening. *Journal of Biotechnology* **324**: 211–219.
- Corpas FJ, González-Gordo S, Palma JM (2021). Nitric oxide and hydrogen sulfide modulate the NADPH-generating enzymatic system in higher plants. *Journal of Experimental Botany* **72**: 830–847.
- Corpas FJ, González-Gordo S, Cañas A, Palma JM (2019). Nitric oxide and hydrogen sulfide in plants: Which comes first? *Journal of Experimental Botany* **70**: 4391–4404.
- Corpas FJ, Palma JM (2020). H₂S signaling in plants and applications in agriculture. *Journal of Advanced Research* **24**: 131–137.
- Fares A, Rossignol M, Peltier JB (2011). Proteomics investigation of endogenous S-nitrosylation in Arabidopsis. *Biochemical and Biophysical Research Communications* **416**: 331–336.
- Filipovic MR, Zivanovic J, Alvarez B, Banerjee R (2018). Chemical biology of H₂S signaling through persulfidation. *Chemical Reviews* **118**: 1253–1337.
- González-Gordo S, Bautista R, Claros MG, Cañas A, Palma JM, Corpas FJ (2019). Nitric oxide-dependent regulation of sweet pepper fruit ripening. *Journal of Experimental Botany* **70**: 4557–4570.
- González-Gordo S, Palma JM, Corpas FJ (2020). Appraisal of H₂S metabolism in *Arabidopsis thaliana*: In silico analysis at the subcellular level. *Plant Physiology and Biochemistry* **155**: 579–588.
- Hancock JT, Whiteman M (2016). Hydrogen sulfide signaling: Interactions with nitric oxide and reactive oxygen species. *Annals of the New York Academy of Sciences* **1365**: 5–14.
- Hughes MN (2008). Chemistry of nitric oxide and related species. *Methods Enzymology* **436**: 3–19.
- Iqbal N, Umar S, Khan NA, Corpas FJ (2021). Nitric oxide and hydrogen sulfide coordinately reduce glucose sensitivity and decrease oxidative stress via ascorbate-glutathione cycle in heat-stressed wheat (*Triticum aestivum* L.) plants. *Antioxidants (Basel)* **10**: 108.
- Jiménez A, Hernández JA, Pastori G, del Río LA, Sevilla F (1998). Role of the ascorbate-glutathione cycle of mitochondria and peroxisomes in the senescence of pea leaves. *Plant Physiology* **118**: 1327–1335.
- Kabil O, Banerjee R (2010). Redox biochemistry of hydrogen sulfide. *Journal of Biology Chemistry* **285**: 21903–21907.
- Kato H, Takemoto D, Kawakita K (2013). Proteomic analysis of S-nitrosylated proteins in potato plant. *Physiologia Plantarum* **148**: 371–386.
- Kharbech O, Sakouhi L, Ben Massoud M, Jose Mur LA, Corpas FJ, Djebali W, Chaoui A (2020). Nitric oxide and hydrogen sulfide protect plasma membrane integrity and mitigate chromium-induced methylglyoxal toxicity in maize seedlings. *Plant Physiology and Biochemistry* **157**: 244–255.
- Kolluru GK, Shen X, Kevil CG (2013). A tale of two gases: NO and H₂S, foes or friends for life? *Redox Biology* **1**: 313–318.
- Kolluru GK, Yuan S, Shen X, Kevil CG (2015). H₂S regulation of nitric oxide metabolism. *Methods Enzymology* **554**: 271–297.
- Lamattina L, García-Mata C, Graziano M, Pagnussat G (2003). Nitric oxide: The versatility of an extensive signal molecule. *Annual Review of Plant Biology* **54**: 109–136.
- Leshem YY, Wills RBH, Veng-Va Ku V (1998). Evidence for the function of the free radical gas-nitric oxide (NO) as an endogenous maturation and senescence regulating factor in higher plants. *Plant Physiology and Biochemistry* **36**: 825–833.
- Lindermayr C, Saalbach G, Durner J (2005). Proteomic identification of S-nitrosylated proteins in Arabidopsis. *Plant Physiology* **137**: 921–930.
- Liu YL, Shen ZJ, Simon M, Li H, Ma DN, Zhu XY, Zheng HL (2019). Comparative proteomic analysis reveals the regulatory effects of H₂S on salt tolerance of mangrove plant *Kandelia obovata*. *International Journal of Molecular Science* **21**: 118.
- Maruta T, Ishikawa T (2018). Ascorbate peroxidase functions in higher plants: The control of the balance between oxidative damage and signaling. In: Gupta DK, Palma JM, Corpas FJ (eds.), *Antioxidants and Antioxidant Enzymes in Higher Plants*, pp. 41–60. Springer.
- Maruta T, Sawa Y, Shigeoka S, Ishikawa T (2016). diversity and evolution of ascorbate peroxidase functions in chloroplasts: More than just a classical antioxidant enzyme? *Plant and Cell Physiology* **57**: 1377–1386.
- Yoshimura K, Yabuta Y, Tamoi M, Ishikawa T, Shigeoka S (1999). Alternatively spliced mRNA variants of chloroplast ascorbate peroxidase isoenzymes in spinach leaves. *Biochemical Journal* **338**: 41–48.
- McCleverty JA (2004). Chemistry of nitric oxide relevant to biology. *Chemical Reviews* **104**: 403–418.
- Mukherjee S, Corpas FJ (2020). Crosstalk among hydrogen sulfide (H₂S), nitric oxide (NO) and carbon monoxide (CO) in root-system development and its rhizosphere interactions: A gaseous interactome. *Plant Physiology and Biochemistry* **155**: 800–814.
- Muñoz-Vargas MA, González-Gordo S, Cañas A, López-Jaramillo J, Palma JM, Corpas FJ (2018). Endogenous hydrogen sulfide (H₂S) is up-regulated during sweet pepper (*Capsicum annuum* L.) fruit ripening. *In vitro* analysis shows that NADP-dependent isocitrate dehydrogenase (ICDH) activity is inhibited by H₂S and NO. *Nitric Oxide* **81**: 36–45.
- Niu L, Yu J, Liao W, Xie J, Yu J, Lv J, Xiao X, Hu L, Wu Y (2019). Proteomic investigation of S-nitrosylated proteins during NO-induced adventitious rooting of cucumber. *International Journal of Molecular Sciences* **20**: 5363.
- Muñoz-Vargas MA, González-Gordo S, Palma JM, Corpas FJ (2020). Inhibition of NADP-malic enzyme activity by H₂S and NO in sweet pepper (*Capsicum annuum* L.) fruits. *Physiologia Plantarum* **168**: 278–288.
- Netto LES, de Oliveira MA, Monteiro G, Demasi APD, Cussiol JRR, Discola KF, Demasi M, Silva GM, Alves SV, Faria VG, Horta BB (2007). Reactive cysteine in proteins: Protein folding, antioxidant defense, redox signaling and more. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* **146**: 180–193.
- Palma JM, Mateos RM, López-Jaramillo J, Rodríguez-Ruiz M, González-Gordo S, Lechuga-Sancho AM, Corpas FJ (2020). Plant catalases as NO and H₂S targets. *Redox Biology* **34**: 101525.

- Palmer RM, Ferrige AG, Moncada S (1987). Nitric oxide release accounts for the biological activity of endothelium-derived relaxing factor. *Nature* **327**: 524–526.
- Ramirez L, Simontacchi M, Murgia I, Zabaleta E, Lamattina L (2011). Nitric oxide, nitrosyl iron complexes, ferritin and frataxin: A well equipped team to preserve plant iron homeostasis. *Plant Science* **181**: 582–592.
- Rodríguez-Ruiz M, Mateos RM, Codesido V, Corpas FJ, Palma JM (2017). Characterization of the galactono-1,4-lactone dehydrogenase from pepper fruits and its modulation in the ascorbate biosynthesis. *Role of nitric oxide. Redox Biology* **12**: 171–181.
- Scuffi D, Álvarez C, Laspina N, Gotor C, Lamattina L, García-Mata C (2014). Hydrogen sulfide generated by L-cysteine desulphydrase acts upstream of nitric oxide to modulate abscisic acid-dependent stomatal closure. *Plant Physiology* **166**: 2065–2076.
- Shigeoka S, Ishikawa T, Tamoi M, Miyagawa Y, Takeda T, Yabuta Y, Yoshimura K (2002). Regulation and function of ascorbate peroxidase isoenzymes. *Journal of Experimental Botany* **53**: 1305–1319.
- Simontacchi M, Jasid S, Puntarulo S (2004). Nitric oxide generation during early germination of Sorghum seeds. *Plant Science* **167**: 839–847.
- Stamler JS, Singel DJ, Loscalzo J (1992). Biochemistry of nitric oxide and its redox-activated forms. *Science* **258**: 18980–18902.
- Tanou G, Job C, Rajjou L, Arc E, Belghazi M, Diamantidis G, Molassiotis A, Job D (2009). Proteomics reveals the overlapping roles of hydrogen peroxide and nitric oxide in the acclimation of citrus plants to salinity. *Plant Journal* **60**: 795–804.
- Wolhuter K, Whitwell HJ, Switzer CH, Burgoyne JR, Timms JF, Eaton P (2018). Evidence against stable protein S-nitrosylation as a widespread mechanism of post-translational regulation. *Molecular Cell* **69**: 438–450.
- Yamaguchi K, Mori H, Nishimura M (1995). A novel isoenzyme of ascorbate peroxidase localized on glyoxysomal and leaf peroxisomal membranes in pumpkin. *Plant and Cell Physiology* **36**: 1157–1162.
- Yang H, Mu J, Chen L, Feng J, Hu J, Li L, Zhou JM, Zuo J (2015). S-nitrosylation positively regulates ascorbate peroxidase activity during plant stress responses. *Plant Physiology* **167**: 1604–1615.
- Zhou ZH, Wang Y, Ye XY, Li ZG (2018). Signaling Molecule Hydrogen sulfide improves seed germination and seedling growth of maize (*Zea mays* L.) under high temperature by inducing antioxidant system and osmolyte biosynthesis. *Front in Plant Science* **9**: 1288.
- Zuccarelli R, Rodríguez-Ruiz M, Lopes-Oliveira PJ, Pascoal GB, Andrade SCS, Furlan CM, Purgatto E, Palma JM, Corpas FJ, Rossi M, Freschi L (2021). Multifaceted roles of nitric oxide in tomato fruit ripening: NO-induced metabolic rewiring and consequences for fruit quality traits. *Journal of Experimental Botany* **72**: 941–958.