

The Inhibitory Effects of HYDAMTIQ, a Novel PARP Inhibitor, on Growth in Human Tumor Cell Lines With Defective DNA Damage Response Pathways

Enrico Mini,* Ida Landini,* Laura Lucarini,† Andrea Lapucci,* Cristina Napoli,‡
Gabriele Perrone,* Renato Tassi,* Emanuela Masini,† Flavio Moroni,† and Stefania Nobili‡

*Department of Experimental and Clinical Medicine, University of Florence, Florence, Italy

†Department of NEUROFARBA, University of Florence, Florence, Italy

‡Department of Health Sciences, University of Florence, Florence, Italy

The poly(ADP-ribose) polymerase (PARP) enzymes play a key role in the regulation of cellular processes (e.g., DNA damage repair, genomic stability). It has been shown that PARP inhibitors (PARPIs) are selectively cytotoxic against cells having dysfunctions in genes involved in DNA repair mechanisms (synthetic lethality). Drug-induced PARP inhibition potentiates the activity of anticancer drugs such as 5-fluorouracil in enhancing DNA damage, whose repair involves PARP-1 activity. The aim of this study was to evaluate the inhibitory effects of a novel PARPI, HYDAMTIQ, on growth in human tumor cell lines characterized by different features with regard to DNA damage response pathways (*BRCA* mutational status, microsatellite status, and ATM expression level) and degree of sensitivity/resistance to 5-fluorouracil. HYDAMTIQ showed a more potent inhibitory effect on cell growth in a *BRCA2* mutant cell line (CAPAN-1) compared with wild-type cells (C2-6, C2-12, and C2-14 CAPAN-1 clones, and MCF-7). No statistically significant difference was observed after HYDAMTIQ exposure between cells having a different MS status or a different MRE11 mutational status. HYDAMTIQ induced greater antiproliferative effects in SW620 cells expressing a low level of ATM than in H630 cells expressing a high level of ATM. Finally, the combination of HYDAMTIQ and 5-fluorouracil exerted a synergistic effect on the inhibition of SW620 cell growth and an antagonistic effect on that of H630 cell growth. Our results show that the novel PARP inhibitor HYDAMTIQ potently inhibits the growth of human tumor cells with defective DNA damage response pathways and exerts synergistic cytotoxicity in combination with 5-fluorouracil. These data provide relevant examples of synthetic lethality and evidence for further development of this novel PARPI.

Key words: PARP inhibitors (PARPIs); HYDAMTIQ; 5-Fluorouracil (5-FU); Human tumor cell lines; DNA damage response

INTRODUCTION

Poly(ADP-ribose) polymerases (PARPs) are a major family of proteins that catalyzes posttranslational modifications [i.e., poly(ADP-ribosyl)ation and mono(ADP-ribosyl)ation (PARylation and MARYlation, respectively)]¹. Such proteins use the NAD⁺ cofactor to transfer an ADP-ribose group onto a specific protein acceptor with the concomitant release of nicotinamide². Today, at least 17 members of this family of enzymes have been identified. PARPs regulate both nuclear and cytoplasmic processes, including DNA damage repair, transcription, cell cycle progression, unfolded protein response, mitosis, and cell death².

The nuclear members (PARP-1–3) have been reported to be involved in several biological processes, including DNA

damage. Since the effects of PARP inhibitors (PARPIs) on cancer cells are related to DNA damage, most research on PARPIs has focused on such PARPs. PARP-1 and PARP-2 synthesize branched PAR polymers, whereas PARP-3 is a critical player in the stabilization of the mitotic spindle and in telomere integrity, notably by associating and regulating the mitotic components NuMA and tankyrase 1³.

Emerging data suggest that PARP inhibition is a potentially important strategy for managing a significant subset of tumors. The rationale for the use of PARPIs is based on the exploitation of synthetic lethality conferred to cancer cells by the copresence of an aberrant DNA damage response (DDR) gene and the PARPI⁴.

To date, only two PARPIs are approved for clinical use: olaparib and rucaparib. Olaparib received accelerated

Address correspondence to Enrico Mini, Department of Experimental and Clinical Medicine, University of Florence, Viale Pieraccini, 6, 50139 Firenze, Italy. Tel: +39-055-2758368; E-mail: enrico.mini@unifi.it or Stefania Nobili, Department of Health Sciences, University of Florence, Viale Pieraccini, 6, 50139 Firenze, Italy. Tel: +39-055-2758386; E-mail: stefania.nobili@unifi.it

approval by the FDA in pretreated ovarian cancer patients harboring germline *BRCA* mutations on the basis of favorable objective response rate and duration of response obtained in clinics⁵. Rucaparib has been recently approved for the treatment of platinum-sensitive *BRCA*-mutated advanced ovarian cancer patients who received more than two lines of platinum-based therapy and whose tumors show homologous recombination (HR) loss of heterozygosity⁶. In cells with mutations in *BRCA1* and/or *BRCA2* that are HR deficient, PARPis stimulate DNA repair by the error-prone nonhomologous end joining recombination (NHEJ) mechanism. Thus, NHEJ is the major contributor to the cytotoxicity observed in HR-deficient cells treated with PARPis⁷.

The PARPI HYDAMTIQ, a hydroxyl derivative of thieno[2,3-*c*]isoquinolin-5(4*H*)-one, is a potent and selective PARP-1/2 inhibitor^{8,9}. The neuroprotective¹⁰, antiasthmatic¹¹, and anti-inflammatory¹² properties of HYDAMTIQ have been investigated in *in vivo* experimental systems in relation to the fact that PARP-1 activation contributes to postischemic brain damage and facilitates the expression of tumor necrosis factors and other proinflammatory cytokines.

The present research investigates the potential anticancer activity of HYDAMTIQ by studying the inhibitory effects of HYDAMTIQ on the growth of a panel of human cancer cell lines characterized by differences in the mutational status of *BRCA* gene, microsatellite status, expression levels of ATM (ataxia-telangiectasia mutated), and degree of intrinsic sensitivity/resistance to 5-fluorouracil (5-FU).

Cell lines received long-term exposure to HYDAMTIQ in order to closely mimic the exposure times that derive from the continuous oral administration of olaparib in clinics^{13,14}.

MATERIALS AND METHODS

Drugs and Supplies

Olaparib was obtained from Selleck Chemicals (Munich, Germany); the novel PARPI HYDAMTIQ was provided by Prof. Pellicciari (University of Perugia) and TESPHERMA (Perugia, Italy). 5-FU and sulforhodamine B (SRB) were obtained from Sigma-Aldrich (Milan, Italy). All the other chemicals were of analytical grade. RPMI-1640 and high-glucose DMEM cell culture mediums, fetal calf serum (FCS), trypsin-EDTA, antibiotics, and phosphate-buffered saline were obtained from Gibco (Milan, Italy).

The phospho-ATM (Ser1981) mouse monoclonal antibody was obtained from GeneTex (Irvine, CA, USA) and Alexa Fluor[®] 488 and 4,6-diamidino-2 phenylindole (DAPI) were from Abcam (Cambridge, UK).

Cell Lines and Culture Conditions

The human *BRCA2*-mutated pancreatic cancer cell line (CAPAN-1) sensitive to cisplatin and its clones (C2-6,

C2-12, and C2-14) characterized by secondary intragenic mutations in *BRCA2* that restore the wild-type *BRCA2* reading frame and that confer resistance to cisplatin were a gift from Dr. Toshiyasu Taniguchi (Fred Hutchinson Cancer Research Center, Seattle, WA, USA). The breast cancer cell line MCF-7 (*BRCA1/2* wild type) and the colorectal cancer cell lines HCT-8, HCT-116, H630, LoVo, DLD-1 [microsatellite instable (MSI)], HT29, and SW620 [microsatellite stable (MSS)] were obtained from ATCC (Manassas, VA, USA). Cell lines were maintained in an appropriate culture medium (RPMI-1640 or high-glucose DMEM), supplemented with 10% FCS and antibiotics (penicillin, 100 U/ml; streptomycin, 100 µg/ml) at 37°C in a 5% CO₂ humidified atmosphere and subcultured twice weekly.

Cell Growth Inhibition Studies

The antiproliferative effects of HYDAMTIQ and olaparib were evaluated on the growth of the studied human tumor cell lines by the SRB assay¹⁵. Briefly, exponentially growing cells were inoculated into 96-well microtiter plates at plating densities of 2×10^3 cells/well. After cell inoculation, the microtiter plates were incubated under standard culture conditions (37°C, 5% CO₂, 95% air, and 100% relative humidity) for 24 h prior to the addition of experimental drugs. After 24 h, the medium was removed and replaced with medium containing PARPis at concentrations ranging from 0.5 to 50 µM for continuous exposures of 72, 144, or 240 h. The assay was completed by the addition of cold TCA. Cells were fixed *in situ* by 10% TCA and stained by SRB solution at 0.4% (w/v) in 1% acetic acid. After staining, unbound dye was removed by washing five times with 1% acetic acid, and the plates were air dried. Bound stain was subsequently solubilized with 10 mM Tris base, and the absorbance was read on an automated plate reader at a wavelength of 540 nm.

The IC₂₀, IC₂₅, and IC₅₀ resulting in a 20%, 25%, and 50% reduction in the net protein content (as measured by SRB staining) in drug-treated cells compared to untreated control cells were determined after drug exposures.

The degree of resistance (*R*) to PARPis is defined as the ratio between IC₅₀ values obtained in *BRCA1/2*-mutated, MSI, or lowest-ATM expression cell lines and the *BRCA1/2* wild type, MSS, and highest-ATM expression cell lines, respectively. *R* values lower than 0.9 indicate collateral sensitivity, and those between 0.9 and 1.1 equal sensitivity and *R* values higher than 1.1 indicate cross-resistance¹⁶.

The growth inhibitory effects of the combination of PARPis and 5-FU on SW620 and H630 have been studied using the IC₂₅ of 5-FU (1 and 3 µM for H630 and SW620 cells, respectively) and the IC₅₀ of HYDAMTIQ (20 and 45 µM for SW620 and H630 cells, respectively)

or of olaparib (5 and 30 μM for SW620 and H630, respectively) and by exposing cells to drugs for 144 h.

The potential synergism between HYDAMTIQ and 5-FU was evaluated using Drewinko's fraction method¹⁶. The synergistic, additive, and antagonistic interactions were determined by the ratio between the cell growth following single-agent exposures (expected value) and the cell growth following drug-combination exposures (observed cell growth) (CI) as follows: $\text{CI} < 0.9$ synergism, CI from 0.9 to 1.1 additivity, and $\text{CI} > 1.1$ antagonism.

Immunofluorescence Analysis

Immunofluorescence analysis of baseline ATM protein levels was performed in duplicate in five human colon carcinoma cell lines (HCT-8, H630, SW620, LoVo, and HT29). By densitometry analysis of protein levels, the cell lines with the highest and lowest level of ATM were identified (H630 and SW620, respectively). Such cell lines were used in the experiments of expression modulation after long-term exposure (240 h) at $2 \times \text{IC}_{50}$ of HYDAMTIQ.

Cells were grown in glass chamber slides at a starting concentration of 0.75×10^4 in 0.5 ml of complete culture medium for 4–5 days until confluence. Slides were washed twice with cold PBS and then fixed with cold 100% methanol for 10 min at -20°C . Cells were then permeabilized with 0.1% Triton X-100 in PBS for 15 min.

After washing three times with PBS, the slides were blocked for 30 min in PBS containing 2% BSA at room temperature, and then incubated for 1 h with 1:1,000 dilution of phospho-ATM(Ser1981) antibody in 2% BSA/PBS. After several washings with 0.1% Tween 20/PBS, bound antibodies were detected with a 1:200 dilution of goat anti-mouse secondary polyclonal antibody, FITC conjugate (Abcam). Nuclei were counterstained with a 1:5,000 dilution of DAPI for 15 min. Representative images were acquired by an Olympus BX63 microscope coupled with a CellSens Dimension Imaging Software version 1.6 (Olympus, Milan, Italy).

The fluorescence intensity was determined in a semi-quantitative way with densitometry analysis (ImageJ software, NIH Image) normalizing for the number of cells per microscope field.

Statistical Analysis

Statistical analysis was performed employing Student's *t*-test and Spearman test (GraphPad Prism v.5). Values of $p < 0.05$ were considered significant.

RESULTS

HYDAMTIQ Inhibitory Effects on the Growth of Tumor Cell Lines With Different Mutational Status of BRCA2

The inhibitory effects of HYDAMTIQ and olaparib were evaluated in the *BRCA2*-mutated CAPAN-1 cell lines, its C2-6, C2-12, and C2-14 clones characterized

by secondary intragenic mutations in *BRCA2* that restore the wild-type *BRCA2* reading frame, and in the *BRCA2* wild-type MCF-7 cells after 72, 144, and 240 h of exposure. Slightly higher IC_{50} values were observed with HYDAMTIQ compared with olaparib after 72 or 144 h of treatment, whereas after 240 h HYDAMTIQ and olaparib showed similar IC_{50} values (Table 1).

At all exposure times, HYDAMTIQ was significantly more active at inhibiting CAPAN-1 cell growth compared to C2-6, C2-12, and C2-14 *BRCA2* wild-type clones. Resistance ratio (*R*) values ranged from 1.8 to 2.1 after 72 and 144 h and from 2.0 to 2.4 after 240 h of exposure. Similarly, HYDAMTIQ was more active in CAPAN-1 cells than in MCF-7 cells. *R* values ranged from 2.7 and 5.6 after all the interval exposure times were tested. Overall, a time-dependent inhibitory effect on cell growth was observed. Lower IC_{50} values were associated with prolonged time of exposure ($p < 0.05$). A similar trend was observed for olaparib ($p < 0.05$) (Table 1). The inhibitory effects of olaparib on cell growth compared to HYDAMTIQ were more marked following 72 and 144 h of treatment, whereas after 240 h the IC_{50} values of the two drugs were similar.

HYDAMTIQ Inhibitory Effects on the Growth of Tumor Cell Lines With Different Status of Microsatellites

The IC_{50} values of HYDAMTIQ and olaparib in the microsatellite stable HT29 and SW630 cell lines (MSS) have been compared with those obtained in the microsatellite unstable HCT-8, HCT-116, LoVo, and DLD-1 cell lines (MSI). No statistically significant difference was observed after 144 (data not shown) or 240 h of exposure to HYDAMTIQ or olaparib between cells with different MS status (Fig. 1). Furthermore, no difference was observed in such cell lines according to the MRE11 mutational status (Fig. 1).

HYDAMTIQ Inhibitory Effects on the Growth of Tumor Cell Lines With Variable Levels of ATM Protein

A panel of colorectal cancer cell lines (HCT-8, H630, SW620, LoVo, and HT29) was analyzed for ATM protein expression levels by immunofluorescence assay. Among the colorectal cancer cell lines, a high degree of variability in ATM protein expression was observed. In particular, H630 cells had the highest level of ATM protein and SW620 cells the lowest (H630 > HCT-8 > HT29 > LoVo > SW620) (Fig. 2).

Relationships between IC_{50} values of HYDAMTIQ or olaparib obtained at 144 and 240 h in the five colorectal cancer cell lines and ATM protein expression levels were investigated. Direct associations were observed for all the comparisons performed, and statistically significant values were obtained between 144-h IC_{50} of HYDAMTIQ and ATM level ($p = 0.0167$) and between 144-h (Fig. 3A and C)

Table 1. Inhibitory Effects of HYDAMTIQ and Olaparib on Cell Growth of Tumor Cell Lines With Different *BRCA1/2* Mutational Status After 72, 144, and 240 h of Exposure

	HYDAMTIQ, IC ₅₀ (μM±SE)			Olaparib, IC ₅₀ (μM±SE)		
	72 h	144 h	240 h	72 h	144 h	240 h
CAPAN-1	16.8±0.3	5.4±0.1	1.9±0.1	9.4±0.8	2.4±0.1	1.9±0.1
<i>n</i>	3	3	3	3	3	3
CAPAN-1 C2-6	30.2±1.0	9.7±0.8	4.3±0.2	13.3±0.8	5.6±0.2	4.4±0.3
<i>r</i>	1.8	1.8	2.3	1.4	2.4	2.4
<i>n</i>	3	3	3	3	3	3
<i>p</i>	0.002	0.037	0.002	0.146	0.0010	0.011
CAPAN-1 C2-12	33.7±1.3	11.6±1.0	4.5±0.1	15.1±0.3	7.7±0.2	4.3±0.1
<i>r</i>	2.0	2.1	2.4	1.6	3.3	2.3
<i>n</i>	3	3	3	3	3	3
<i>p</i>	0.002	0.023	0.001	0.016	0.0001	0.0006
CAPAN-1 C2-14	34.5±1.0	9.7±0.4	3.9±0.1	18.5±2.0	7.6±0.2	3.1±0.2
<i>r</i>	2.1	1.8	2.0	1.5	2.9	1.8
<i>n</i>	3	3	3	3	3	3
<i>p</i>	0.001	0.004	0.001	0.004	<0.0001	0.042
MCF-7	44.7±2.6	30.4±3.4	8.3±0.9	26.1±0.6	19.9±0.9	3.3±0.3
<i>r</i>	2.7	5.6	4.4	2.8	8.4	1.8
<i>n</i>	4	4	3	4	4	3
<i>p</i>	0.007	0.026	0.026	0.003	0.004	0.054

IC₅₀, concentration of drug required to inhibit cell growth by 50%; SE, standard error; *r*, resistance index; *n*, number of experiments; *p*, HYDAMTIQ or olaparib IC₅₀ values of CAPAN-1 cells versus CAPAN-1 clones or MCF-7 cells.

or 240-h (Fig. 3B and D) IC₅₀ of olaparib and ATM level ($p=0.0167$ in both cases).

Different sensitivities of colorectal cancer cells to PARPIs according to ATM levels is shown in Figure 4. Resistance values given by the ratio of the highest ATM expression cells (H630) over the lower ATM expression cells (SW620) after 240 h of treatment with HYDAMTIQ or olaparib were 0.5 and 0.3, respectively (Fig. 4A and B).

After 144 h of treatment with HYDAMTIQ or olaparib, the *r* values according to the same parameters were 0.4 and 0.3, respectively (data not shown).

In order to investigate the relationships between PARP inhibition and ATM protein expression variability, H630 and SW620 cells were exposed to a double IC₅₀ concentration of HYDAMTIQ for 240 h. Comparison of ATM expression levels before and after exposure to




Figure 1. Inhibition of human colon carcinoma cell growth: dose-dependent effects of HYDAMTIQ (A) and olaparib (B). Black curves indicate microsatellite stable (MSS) cell lines [*MRE11* wild type (wt)]; blue curves indicate microsatellite unstable (MSI) cell lines with heterozygous status of *MRE11*; and red curves indicate MSI cell lines with homozygous mutational status of *MRE11*. Each curve indicates the concentration–cell growth inhibition curve of a single cell line following 240 h of exposure.




Figure 2. Baseline levels of ATM (ataxia-telangiectasia mutated) protein expression in colon carcinoma cell lines. (A) Immunofluorescence staining of cell lines labeled with anti-ATM polyclonal antibody (green) and nuclei (blue) counterstained with DAPI (4,6-diamidino-2 phenylindole) (40 \times). Images show the baseline levels of ATM in cell lines. ATM plays a critical role in signaling DNA damage to cell cycle checkpoints and DNA repair pathways. (B) Baseline ATM protein level optical density according to the different cell lines tested. *au, arbitrary units, mean \pm standard error.

HYDAMTIQ showed a decrease, although not statistically significant, of intranuclear fluorescent ATM protein in both cell lines (Fig. 4C and D).

Inhibitory Effects of Combined HYDAMTIQ With 5-Fluorouracil on Tumor Cell Lines With High or Low ATM Protein Levels

H630 and SW620 cells were exposed to 5-FU combined with HYDAMTIQ or olaparib for 144 h. Cell growth inhibition was evaluated using H630 and SW620 IC₂₅ values for 5-FU and IC₅₀ values for HYDAMTIQ or olaparib.

There was a synergistic effect obtained following exposure to both combinations (i.e., HYDAMTIQ/5-FU or olaparib/5-FU, CI: 0.6 or 0.5, respectively) in SW620 cells, characterized by low ATM level (Fig. 5A and B). On the contrary, according to the same experimental conditions, an antagonistic effect was observed in H630 cells characterized by high ATM level (CI: 1.3 and 1.6, respectively) (Fig. 5C and D).

DISCUSSION

PARPs are a superfamily of enzymes involved in the regulation of cellular processes, such as DNA repair, gene transcription, cell cycle progression, cell death and genomic stability, and apoptosis^{1,17,18}.

PARP-1 and PARP-2 control single-strand break (SSB) repair systems such as base excision repair (BER) and nucleotide excision repair (NER) pathways¹⁹. PARP-1 is also involved in the transcription of mitochondrial proteins²⁰. In addition, PARP-1, PARP-2, and PARP-3 are

involved in double-stranded break (DSB) repair. In particular, PARP-3 has been shown to be involved in DSB repair, mainly promoting the NHEJ process¹⁹. DNA damage repair systems, together with cell cycle checkpoints, are the cellular reaction to exogenous and endogenous genotoxic injuries that may produce DNA SSBs and DNA DSBs. Overall, DDR plays a pivotal role in maintaining genomic integrity and avoiding the occurrence of genomic instability typical of cancer cells. However, several examples of aberrant proteins are known today [e.g., MRE11 in DSB repair, BRCA1 and BRCA2 in HR repair (HRR), and ATM and ATR (i.e., ATM and RAD3 related) in cell cycle checkpoints]^{21,22}. ATM, ATR, and DNA-dependent protein kinases (DNA-PKcs) are the most upstream DDR kinases. In response to DNA damage, several proteins are phosphorylated in an ATM- or ATR-dependent manner, whereas DNA-PKcs regulate a smaller number of targets and play a role, primarily in NHEJ²³.

Most PARPIs act as competitive inhibitors of NAD by interfering at the binding site of the enzyme²⁴ or trapping PARP enzymes at damaged DNA²⁵. In the last few years, a significantly increased potency and selectivity of PARPIs have been obtained because of the advancements in the knowledge of relationships between chemical structures and pharmacological activity^{26,27}.

We investigated the antiproliferative properties of the PARPI HYDAMTIQ on the growth of a panel of human tumor cell lines. HYDAMTIQ is a potent inhibitor of PARP-1 and PARP-2 with IC₅₀ values in the nanomolar range (IC₅₀ 29 and 38 nM, respectively)⁹. The biological




Figure 3. Correlations between baseline ATM expression and HYDAMTIQ IC_{50} values in colon carcinoma cell lines. Scatter plots indicate relationships between baseline ATM expression and IC_{50} of HYDAMTIQ at 144 (A) and 240 (B) h or IC_{50} of olaparib at 144 (C) and 240 (D) h in cell lines. *au, arbitrary units, mean \pm standard error. Black circles indicate the intersection of IC_{50} /ATM expression values of a single cell line.

activity of HYDAMTIQ has been so far investigated in vivo inflammatory models only^{10–12}.

Our study showed that HYDAMTIQ and olaparib were more active in the *BRCA2*-mutated CAPAN-1 human pancreatic cancer cell line than in the wild-type *BRCA2*, cisplatin-resistant CAPAN-1 clones (i.e., C2-6, C2-12, and C2-14) and in the *BRCA1/2* wild-type MCF-7 breast cancer cell line. No difference in the activity of HYDAMTIQ or olaparib was observed in cell lines with different *MS* or *MRE11* status.

HYDAMTIQ and olaparib were more active in SW620 human colorectal cancer cells characterized by a low ATM protein expression than in H630 human colorectal cancer cells with high ATM protein levels. Following exposure to HYDAMTIQ, ATM protein levels decreased in both study cell lines, although a more marked decrease was observed

in SW620 cells. In the same low- and high-ATM expression cell lines, the potential synergistic effect of the combination of 5-FU and HYDAMTIQ or olaparib was evaluated. In SW620 cells, the combination of 5-FU and HYDAMTIQ or olaparib exerted a synergistic effect, whereas an antagonistic effect in H630 cells was observed.

The major determinant of the PARPI activity is the presence of defects in the DNA repair mechanisms of cancer cells. In particular, PARPIs are selectively cytotoxic against cells with mutations of genes involved in DNA repair (synthetic lethality)^{21,22,28}. The absence of defects in DNA repair mechanisms does not allow an optimal inhibitory effect on the growth of PARPIs. Ovarian, breast, and pancreatic cancers with mutations in *BRCA1* or *BRCA2* genes involved in the HRR system have been shown to be sensitive to the activity of PARPIs both at preclinical




Figure 4. Dose-dependent effects of HYDAMTIQ and olaparib in SW620 and H630 cell lines and changes of ATM levels following $2 \times \text{IC}_{50}$ HYDAMTIQ treatment. Inhibition of cell growth: dose-dependent effects of HYDAMTIQ (A) or olaparib (B) in cell lines with the highest (H630) or the lowest (SW620) level of ATM protein. Inhibitory cell growth effects were evaluated after 240 h of exposure. ATM protein expression levels in H630 (C) and SW620 (D) cells exposed to HYDAMTIQ $2 \times \text{IC}_{50}$ for 240 h. *au, arbitrary units, mean \pm standard error.

and clinical levels²⁹. As expected, our results showed a more potent activity of HYDAMTIQ in *BRCA2*-mutated cells than in *BRCA* wild-type cells and are in agreement with those reported by Sakai et al.³⁰ concerning the PARPI AG014361 in the same in vitro tumor models. Other authors also reported a higher activity of different PARPIs, such as talazoparib³¹ or AG014699³², in CAPAN-1 cells or other *BRCA1/2*-mutated or epigenetically silenced *BRCA1/2* cell lines compared to *BRCA1/2* wild-type cell lines.

PARPIs can also contribute to synthetic lethality in MSI cancer cells. MSI is a molecular phenotype that arises from defects in the DNA mismatch repair (MMR) system. Loss of MMR function is due to somatic or germline

epigenetic/genetic alterations of *MMR* genes. About 15% of colorectal cancers show MSI due to epigenetic silencing of the *MLH1* gene or to germline mutations in mismatch repair genes (e.g., *MLH1*, *MSH2*, *MSH6*)³³. Other mutations that contribute to the MSI phenotype may be present in oncogenes and tumor suppressor genes, in particular *BRAF*³⁴, *MRE11* (ATM)^{35,36}, and *KRAS*³⁷.

Our findings, showing no difference in the activity of HYDAMTIQ or olaparib in colorectal cancer cell lines as a function of both MS and *MRE11* status, are in agreement with those obtained by Genther Williams et al.³⁸, who observed no difference in the sensitivity to niraparib between MSI-H and MSS colorectal cancer cell lines.




Figure 5. Pharmacodynamic interactions between HYDAMTIQ or olaparib and 5-fluorouracil in SW620 and H630 cell lines. Synergistic inhibitory effects of HYDAMTIQ 20 μ M (\sim IC₅₀) (white bar), 5-FU 3 μ M (\sim IC₂₅) (black bar), HYDAMTIQ 20 μ M plus 5-FU 3 μ M (gray bar) (A) or olaparib 5 μ M (\sim IC₅₀) (white bar), 5-FU 3 μ M (\sim IC₂₅) (black bar), and olaparib 5 μ M plus 5-FU 3 μ M (gray bar) (B) on the growth of SW620 cells characterized by low levels of ATM after 144 h of exposure. Antagonistic inhibitory effects of HYDAMTIQ 45 μ M (\sim IC₅₀) (white bar), 5-FU 1 μ M (\sim IC₂₅) (black bar), HYDAMTIQ 45 μ M plus 5-FU 1 μ M (gray bar) (C) or olaparib 30 μ M (\sim IC₅₀) (white bar), 5-FU 1 μ M (\sim IC₂₅) (black bar), and olaparib 30 μ M plus 5-FU 1 μ M (gray bar) (D) on the growth of H630 cells characterized by high levels of ATM after 144 h of exposure. Data from a representative experiment. The dotted line indicates expected growth. CI, observed/expected growth ratio.

Similar findings were observed by McPherson et al.³⁶ in panels of colorectal cancer cell lines with different MS status treated with the PARPI LT-626. Conversely, Vilar et al.³⁵ showed a higher sensitivity to the PARPI veliparib in MSI *MRE11* mutated colorectal cancer cells compared with MSS *MRE11* wild-type colorectal cancer cells. Differences observed between our results and those of Vilar et al. could be due to the different colorectal cancer cells used. In fact, only four out of their eight cell lines were included in our seven cell line panel.

The concept of “synthetic lethality” may also be extended to ATM that is frequently altered or deleted both in solid and hematologic malignancies. Heterozygous

germline mutations in *ATM* are associated with leukemia, breast, and pancreatic cancers³⁹. *ATM* is a serine threonine kinase that represents a critical component of the early response to DNA damage and activation of cell cycle checkpoints⁴⁰. *ATM*, together with *ATR*, collaborates with the checkpoint proteins Chk2 and Chk1, respectively, to arrest the cell cycle and allow DNA repair. Several studies have shown that *ATM* deficiency or *ATM* harboring mutations confer sensitivity to olaparib in in vitro lymphoid tumor models^{41–43} and in breast⁴⁴, gastric⁴⁵, and colorectal⁴⁶ cancer cells. The HYDAMTIQ activity results in SW620 and H630 cells, selected from a panel of human colorectal cancer cell lines on the basis of their *ATM* protein

expression levels (i.e., the lowest and highest, respectively), are in agreement with data from the studies reported above, performed in *in vitro* models treated by olaparib^{41,43–46}. To investigate a potential differential role for ATM protein levels according to HYDAMTIQ treatment, we evaluated variations in ATM protein levels in SW620 and H630 cells pre- and posttreatment. We observed a more marked decrease in posttreatment ATM in SW620 cells, suggesting that other molecular mechanisms, in addition to ATM, are likely involved in resistance to HYDAMTIQ.

In addition to the above-reported conditions, mainly represented by defects in DNA repair pathways and DNA damage recognition, signaling, and checkpoints system that allow PARPis to exploit synthetic lethality, this phenomenon may also be generated by the concomitant presence of PARPis and DNA-damaging agents. In fact, it has been observed that the suppression of PARP activity increases cell susceptibility to DNA-damaging agents and inhibits strand break rejoining. Thus, the inhibition of PARP could represent an interesting pharmacological strategy in order to enhance the effectiveness of DNA damage induced, for instance, by 5-FU, a pyrimidine antimetabolite that is the backbone of the treatment of colorectal cancer both in the adjuvant and metastatic setting⁴⁷. In particular, 5-FU determines imbalances in deoxynucleotide triphosphate levels as well as the accumulation of uracil and 5-FU at the genomic level. Such events are able to activate the ATR- and ATM-dependent checkpoint signaling pathways and the BER pathway.

Although the ATR- and ATM-dependent checkpoint signaling pathways are activated by 5-FU, their role in the survival of cancer cells treated with this drug is still unclear⁴⁸. The observed synergistic effect of the combination of 5-FU and HYDAMTIQ or olaparib in SW620 cells and the observed antagonistic effect in H630 cells support the hypothesis that low ATM expression may be associated with impaired DNA repair while high ATM expression may not be.

To date, few articles have investigated the effects of *in vitro* combinations, including PARPis and 5-FU, with controversial results^{48–52}. Sakogawa et al.⁵⁰ studied the histone variant γ H2AX as a marker of DSBs and found that combined treatment with 5-FU and the PARPI 3-aminobenzamide (3-AB) induced γ H2AX focus formation with similar increases and kinetics as the single treatment with 5-FU in esophageal cancer cells. These authors then suggested that, in combination with 5-FU, PARPis would not be able to increase DSB. On the contrary, Falzacappa et al.⁵² showed that PARP inhibition by rucaparib enhances the chemosensitization of both myeloid (OCI-AML2) and lymphoid (RPMI-8402) leukemia cells to 5-FU exposure, suggesting a synergistic effect of the combined therapy in killing leukemia cells. Based on the above observations, further studies are therefore

needed to clarify the relationship between ATM, 5-FU, and PARPis.

The results of our *in vitro* studies have shown that the novel PARPI HYDAMTIQ exhibits potent cell growth inhibitory activity against a limited number of tumor cell lines with relevant defects in DDR pathways. In these *in vitro* models, HYDAMTIQ showed comparable antiproliferative effects to olaparib.

The spectrum of antiproliferative activity of HYDAMTIQ remains to be determined on a wider panel of tumor types with particular reference to human cell lines with defects in other HRR proteins recruited by PARP (e.g., XRCC1, topoisomerase I). Thus, a molecular target screen of HYDAMTIQ in a large cell line set with known mutational profile of genes involved in the DNA repair process is warranted and will allow more informed interpretation of HYDAMTIQ spectrum of activity.

These results could be of particular clinical relevance since genetic/epigenetic alterations of these genes have been observed in several types of tumors, providing a basis for preclinical and potential clinical development of PARP inhibition in tumors other than ovarian and breast cancers, such as prostate and pancreatic tumors, and glioblastoma multiforme⁵³.

Also, pharmacokinetics/pharmacodynamics (PK/PD) and maximum tolerated dose (MTD) studies for HYDAMTIQ screening in selected preclinical *in vivo* cancer models are warranted prior to first-in-human trials in patients with mixed solid, refractory, metastatic tumors and further potential proof-of-concept trials in the chosen tumor settings. It is conceivable that these studies are feasible since HYDAMTIQ displays favorable physicochemical and PK properties, in particular a good brain penetration supporting, for instance, potential application in glioblastoma (Pellicciari, unpublished data).

Clinical development of PARPis also comprises the approach of combining PARPis with DNA-damaging agents to achieve chemosensitization. This is based on extensive preclinical studies showing that PARPis enhance the action of temozolomide, cisplatin, topoisomerase I inhibitors, and ionizing radiation in tumor cell lines *in vitro* and in mouse xenograft models of human cancers. Our study also provides evidence of synergism of HYDAMTIQ and a fluoropyrimidine, 5-FU. These findings suggest that combining a fluoropyrimidine and a PARPI may be an innovative therapeutic strategy for colorectal cancer. Phase I and II trials of various PARPis in combination with DNA-damaging agents are ongoing.

In this scenario, it will be interesting to see whether HYDAMTIQ is equivalent or superior to the various PARP-1/2 inhibitors currently in clinical development as either single-agent or combination therapy. Thus, further basic studies of this novel potent PARPI are warranted and are likely to be informative and worthwhile.

ACKNOWLEDGMENTS: *This study was supported by contributions from the Associazione Italiana per la Ricerca sul Cancro (IG No. 15565), Milan (to E. Mini), Ente Cassa di Risparmio di Firenze, Florence (No. 2009/0767 to E. Masini and F. Moroni, No. 2012/0744 to E. Masini, No. 2014/0706 to E. Mini, and No. 2014/0969 to S. Nobili), Associazione Giacomo Onlus, Castiglioncello (to E. Mini), and Gruppo Oncologico Chirurgico Cooperativo Italiano, Florence (to E. Mini).*

REFERENCES

- Kraus WL. PARPs and ADP-ribosylation: 50 years...and counting. *Mol Cell* 2015;58(6):902–10.
- Barkauskaite E, Jankevicius G, Ahel I. Structures and mechanisms of enzymes employed in the synthesis and degradation of PARP-dependent protein ADP-ribosylation. *Mol Cell* 2015;58(6):935–46.
- Boehler C, Gauthier LR, Mortusewicz O, Biard DS, Saliou JM, Bresson A, Sanglier-Cianferani S, Smith S, Schreiber V, Boussin F, Dantzer F. Poly(ADP-ribose) polymerase 3 (PARP3), a newcomer in cellular response to DNA damage and mitotic progression. *Proc Natl Acad Sci USA* 2011;108(7):2783–8.
- Cerrato A, Morra F, Celetti A. Use of poly ADP-ribose polymerase [PARP] inhibitors in cancer cells bearing DDR defects: The rationale for their inclusion in the clinic. *J Exp Clin Cancer Res.* 2016;35(1):179.
- Kaufman B, Shapira-Frommer R, Schmutzler RK, Audeh MW, Friedlander M, Balmaña J, Mitchell G, Fried G, Stemmer SM, Hubert A, Rosengarten O, Steiner M, Loman N, Bowen K, Fielding A, Domchek SM. Olaparib monotherapy in patients with advanced cancer and a germline BRCA1/2 mutation. *J Clin Oncol.* 2015;33(3):244–50.
- Swisher EM, Lin KK, Oza AM, Scott CL, Giordano H, Sun J, Konecny GE, Coleman RL, Tinker AV, O'Malley DM, Kristeleit RS, Ma L, Bell-McGuinn KM, Brenton JD, Cragun JM, Oaknin A, Ray-Coquard I, Harrell MI, Mann E, Kaufmann SH, Floquet A, Leary A, Harding TC, Goble S, Maloney L, Isaacson J, Allen AR, Rolfe L, Yelensky R, Raponi M, McNeish IA. Rucaparib in relapsed, platinum-sensitive high-grade ovarian carcinoma (ARIEL2 Part 1): An international, multicentre, open-label, phase 2 trial. *Lancet Oncol.* 2017;18(1):75–87.
- Patel AG, Sarkaria JN, Kaufmann SH. Nonhomologous end joining drives poly(ADP-ribose) polymerase (PARP) inhibitor lethality in homologous recombination-deficient cells. *Proc Natl Acad Sci USA* 2011;108(8):3406–11.
- Moroni F, Formentini L, Gerace E, Camaioni E, Pellegrini-Giampietro DE, Chiarugi A, Pellicciari R. Selective PARP-2 inhibitors increase apoptosis in hippocampal slices but protect cortical cells in models of post-ischaemic brain damage. *Br J Pharmacol.* 2009;157(5):854–62.
- Pellicciari R, Camaioni E, Gilbert AM, Macchiarulo A, Bikker JA, Shah F, Bard J, Costantino G, Gioiello A, Robertson GM, Sabbatini P, Venturoni F, Liscio P, Carotti A, Bellocchi D, Cozzi A, Wood A, Gonzales C, Zaleska MM, Ellingboe JW, Moroni F. Discovery and characterization of novel potent PARP-1 inhibitors endowed with neuroprotective properties: From TIQ-A to HYDAMTIQ. *Med Chem Commun.* 2011;2:559–65.
- Moroni F, Cozzi A, Chiarugi A, Formentini L, Camaioni E, Pellegrini-Giampietro DE, Chen Y, Liang S, Zaleska MM, Gonzales C, Wood A, Pellicciari R. Long-lasting neuroprotection and neurological improvement in stroke models with new, potent and brain permeable inhibitors of poly(ADP-ribose) polymerase. *Br J Pharmacol.* 2012;165(5):1487–500.
- Lucarini L, Pini A, Gerace E, Pellicciari R, Masini E, Moroni F. Poly(ADP-ribose) polymerase inhibition with HYDAMTIQ reduces allergen-induced asthma-like reaction, bronchial hyper-reactivity and airway remodelling. *J Cell Mol Med.* 2014;18(3):468–79.
- Lucarini L, Durante M, Lanzi C, Pini A, Boccalini G, Calosi L, Moroni F, Masini E, Mannaioni G. HYDAMTIQ, a selective PARP-1 inhibitor, improves bleomycin-induced lung fibrosis by dampening the TGF- β /SMAD signalling pathway. *J Cell Mol Med.* 2017;21(2):324–35.
- Tutt A, Robson M, Garber JE, Domchek SM, Audeh MW, Weitzel JN, Friedlander M, Arun B, Loman N, Schmutzler RK, Wardley A, Mitchell G, Earl H, Wickens M, Carmichael J. Oral poly(ADP-ribose) polymerase inhibitor olaparib in patients with BRCA1 or BRCA2 mutations and advanced breast cancer: A proof-of-concept trial. *Lancet* 2010;376(9737):235–44.
- Audeh MW, Carmichael J, Penson RT, Friedlander M, Powell B, Bell-McGuinn KM, Scott C, Weitzel JN, Oaknin A, Loman N, Lu K, Schmutzler RK, Matulonis U, Wickens M, Tutt A. Oral poly(ADP-ribose) polymerase inhibitor olaparib in patients with BRCA1 or BRCA2 mutations and recurrent ovarian cancer: A proof-of-concept trial. *Lancet* 2010;376(9737):245–51.
- Skehan P, Storeng R, Scudiero D, Monks A, McMahon J, Vistica D, Warren JT, Bokesch H, Kenney S, Boyd MR. New colorimetric cytotoxicity assay for anticancer-drug screening. *J Natl Cancer Inst.* 1990;82(13):1107–12.
- Drewinko B, Loo TL, Brown B, Gottlieb JA, Freireich EJ. Combination chemotherapy in vitro with adriamycin. Observations of additive, antagonistic, and synergistic effects when used in two-drug combinations on cultured human lymphoma cells. *Cancer Biochem Biophys.* 1976;1(4):187–95.
- Schreiber V, Dantzer F, Ame JC, de Murcia G. Poly(ADP-ribose): Novel functions for an old molecule. *Nat Rev Mol Cell Biol.* 2006;7(7):517–28.
- Li M, Yu X. The role of poly(ADP-ribosyl)ation in DNA damage response and cancer chemotherapy. *Oncogene* 2015;34(26):3349–56.
- De Vos M, Schreiber V, Dantzer F. The diverse roles and clinical relevance of PARPs in DNA damage repair: Current state of the art. *Biochem Pharmacol.* 2012;84(2):137–46.
- Lapucci A, Pittelli M, Rapizzi E, Felici R, Moroni F, Chiarugi A. Poly(ADP-ribose) polymerase-1 is a nuclear epigenetic regulator of mitochondrial DNA repair and transcription. *Mol Pharmacol.* 2011;79(6):932–40.
- Curtin NJ. DNA repair dysregulation from cancer driver to therapeutic target. *Nat Rev Cancer* 2012;12(12):801–17.
- O'Connor MJ. Targeting the DNA damage response in cancer. *Mol Cell* 2015;60(4):547–60.
- Maréchal A, Zou L. DNA damage sensing by the ATM and ATR kinases. *Cold Spring Harb Perspect Biol.* 2013;5(9):pii:a012716.
- Ruf A, de Murcia G, Schulz. Inhibitor and NAD⁺ binding to poly(ADP-ribose) polymerase as derived from crystal structures and homology modeling. *Biochemistry* 1998;37(11):3893–900.
- Murai J, Huang SY, Das BB, Renaud A, Zhang Y, Doroshov JH, Ji J, Takeda S, Pommier Y. Trapping of

- PARP1 and PARP2 by clinical PARP inhibitors. *Cancer Res.* 2012;72(21):5588–99.
26. Liscio P, Camaioni E, Carotti A, Pellicciari R, Macchiarulo A. From polypharmacology to target specificity: The case of PARP inhibitors. *Curr Top Med Chem.* 2013;13(23):2939–54.
 27. Thompson JM, Nguyen QH, Singh M, Razorenova OV. Approaches to identifying synthetic lethal interactions in cancer. *Yale J Biol Med.* 2015;88(2):145–55.
 28. Lord CJ, Tutt AN, Ashworth A. Synthetic lethality and cancer therapy: Lessons learned from the development of PARP inhibitors. *Annu Rev Med.* 2015;66:455–70.
 29. Scott CL, Swisher EM, Kaufmann SH. Poly (ADP-ribose) polymerase inhibitors: Recent advances and future development. *J Clin Oncol.* 2015;33(12):1397–406.
 30. Sakai W, Swisher EM, Karlan BY, Agarwal MK, Higgins J, Friedman C, Villegas E, Jacquemont C, Farrugia DJ, Couch FJ, Urban N, Taniguchi T. Secondary mutations as a mechanism of cisplatin resistance in BRCA2-mutated cancers. *Nature* 2008;451(7182):1116–20.
 31. Wang B, Chu D, Feng Y, Shen Y, Aoyagi-Scharber M, Post LE. Discovery and characterization of (8S,9R)-5-fluoro-8-(4-fluorophenyl)-9-(1-methyl-1H-1,2,4-triazol-5-yl)-2,7,8,9-tetrahydro-3H-pyrido[4,3,2-de]phthalazin-3-one (BMN 673, Talazoparib), a novel, highly potent, and orally efficacious poly(ADP-ribose) polymerase-1/2 Inhibitor, as an anticancer agent. *J Med Chem.* 2016;59(1):335–57.
 32. Drew Y, Mulligan EA, Vong WT, Thomas HD, Kahn S, Kyle S, Mukhopadhyay A, Los G, Hostomsky Z, Plummer ER, Edmondson RJ, Curtin NJ. Therapeutic potential of poly(ADP-ribose) polymerase inhibitor AG014699 in human cancers with mutated or methylated BRCA1 or BRCA2. *J Natl Cancer Inst.* 2011;103(4):334–46.
 33. Gelsomino F, Barbolini M, Spallanzani A, Pugliese G, Cascinu S. The evolving role of microsatellite instability in colorectal cancer: A review. *Cancer Treat Rev.* 2016;51:19–26.
 34. Gallois C, Laurent-Puig P, Taieb J. Methylator phenotype in colorectal cancer: A prognostic factor or not? *Crit Rev Oncol Hematol.* 2016;99:74–80.
 35. Vilar E, Bartnik CM, Stenzel SL, Raskin L, Ahn J, Moreno V, Mukherjee B, Iniesta MD, Morgan MA, Rennert G, Gruber SB. MRE11 deficiency increases sensitivity to poly(ADP-ribose)polymerase inhibition in microsatellite unstable colorectal cancers. *Cancer Res.* 2011;71:2632–42.
 36. McPherson LA, Shen Y, Ford JM. Poly (ADP-ribose) polymerase inhibitor LT-626: Sensitivity correlates with MRE11 mutations and synergizes with platinum and irinotecan in colorectal cancer cells. *Cancer Lett.* 2014;343(2):217–23.
 37. Lin EI, Tseng LH, Gocke CD, Reil S, Le DT, Azad NS, Eshleman JR. Mutational profiling of colorectal cancers with microsatellite instability. *Oncotarget* 2015;6(39):42334–44.
 38. Genter Williams SM, Kuznicki AM, Andrade P, Dolinski BM, Elbi C, O'Hagan RC, Toniatti C. Treatment with the PARP inhibitor, niraparib, sensitizes colorectal cancer cell lines to irinotecan regardless of MSI/MSS status. *Cancer Cell Int.* 2015;15(1):14.
 39. Thompson D, Duedal S, Kirner J, McGuffog L, Last J, Reiman A, Byrd P, Taylor M, Easton DF. Cancer risks and mortality in heterozygous ATM mutation carriers. *J Natl Cancer Inst.* 2005;97(11):813–22.
 40. Shiloh Y, Ziv Y. The ATM protein kinase: Regulating the cellular response to genotoxic stress, and more. *Nat Rev Mol Cell Biol.* 2013;14:197–210.
 41. Williamson CT, Muzik H, Turhan AG, Zamò A, O'Connor MJ, Bebb DG, Lees-Miller SP. ATM deficiency sensitizes mantle cell lymphoma cells to poly(ADP-ribose) polymerase-1 inhibitors. *Mol Cancer Ther.* 2010;9(2):347–57.
 42. Weston VJ, Oldreive CE, Skowronska A, Oscier DG, Pratt G, Dyer MJ, Smith G, Powell JE, Rudzki Z, Kearns P, Moss PA, Taylor AM, Stankovic T. The PARP inhibitor olaparib induces significant killing of ATM-deficient lymphoid tumor cells in vitro and in vivo. *Blood* 2010;116(22):4578–87.
 43. Williamson CT, Kubota E, Hamill JD, Klimowicz A, Ye R, Muzik H, Dean M, Tu LR, Gilley D, Magliocco AM, McKay BC, Bebb DG, Lees-Miller SP. Enhanced cytotoxicity of PARP inhibition in mantle cell lymphoma harbouring mutations in both ATM and p53. *EMBO Mol Med.* 2012;4(6):515–27.
 44. Gilardini Montani MS, Prodosmo A, Stagni V, Merli D, Monteonofrio L, Gatti V, Gentileschi MP, Barilà D, Soddu S. ATM-depletion in breast cancer cells confers sensitivity to PARP inhibition. *J Exp Clin Cancer Res.* 2013;32:95.
 45. Kubota E, Williamson CT, Ye R, Elegbede A, Peterson L, Lees-Miller SP, Bebb DG. Low ATM protein expression and depletion of p53 correlates with olaparib sensitivity in gastric cancer cell lines. *Cell Cycle* 2014;13(13):2129–37.
 46. Wang C, Jette N, Moussienko D, Bebb DG, Lees-Miller SP. ATM-deficient colorectal cancer cells are sensitive to the PARP inhibitor olaparib. *Transl Oncol.* 2017;10(2):190–6.
 47. Pettersen HS, Visnes T, Vågbo CB, Svaasand EK, Doseth B, Slupphaug G, Kavli B, Krokan HE. UNG-initiated base excision repair is the major repair route for 5-fluorouracil in DNA, but 5-fluorouracil cytotoxicity depends mainly on RNA incorporation. *Nucleic Acids Res.* 2011;39(19):8430–44.
 48. Huehls AM, Wagner JM, Huntoon CJ, Geng L, Erlichman C, Patel AG, Kaufmann SH, Karnitz LM. Poly(ADP-Ribose) polymerase inhibition synergizes with 5-fluorodeoxyuridine but not 5-fluorouracil in ovarian cancer cells. *Cancer Res.* 2011;71(14):4944–54.
 49. Geng L, Huehls AM, Wagner JM, Huntoon CJ, Karnitz LM. Checkpoint signaling, base excision repair, and PARP promote survival of colon cancer cells treated with 5-fluorodeoxyuridine but not 5-fluorouracil. *PLoS One* 2011;6(12):e28862.
 50. Sakogawa K, Aoki Y, Misumi K, Hamai Y, Emi M, Hihara J, Shi L, Kono K, Horikoshi Y, Sun J, Ikura T, Okada M, Tashiro S. Involvement of homologous recombination in the synergism between cisplatin and poly (ADP-ribose) polymerase inhibition. *Cancer Sci.* 2013;104(12):1593–9.
 51. Shekh K, Khan S, Jena G, Kansara BR, Kushwaha S. 3-Aminobenzamide—A PARP inhibitor enhances the sensitivity of peripheral blood micronucleus and comet assays in mice. *Toxicol Mech Methods* 2014;24(5):332–41.
 52. Falzacappa MV, Ronchini C, Faretta M, Iacobucci I, Di Rorà AG, Martinelli G, Meyer LH, Debatin KM, Orecchioni S, Bertolini F, Pelicci PG. The combination of the PARP inhibitor rucaparib and 5FU is an effective strategy for treating acute leukemias. *Mol Cancer Ther.* 2015;14(4):889–98.
 53. Lord CJ, Ashworth A. BRCAness revisited. *Nat Rev Cancer* 2016;16(2):110–20.